arXiv:1508.02938v1 [math.AP] 12 Aug 2015

Uniqueness of Solution of the Unsteady Filtration Problem in Heterogeneous Porous Media

A. Lyaghfouri[1] and E. Zaouche[2]
[1] American University of Ras Al Khaimah
Ras Al Khaimah, UAE
[2] Ecole Normale Supérieure
Algiers, Algeria

Abstract

We establish uniqueness of the solution of the unsteady state dam problem in the heterogeneous and rectangular case assuming the dam wet at the bottom and dry near to the top.

2010 Mathematics Subject Classification: 35A02; 35R35; 76S05.

Key words: Unsteady state dam problem; Fluid flow; Heterogeneous porous medium; Uniqueness of solution.

1 Introduction

We consider a heterogeneous porous medium supplied by several reservoirs of a fluid, represented by a bounded domain Ω of $\mathbb{R}^n$ with locally Lipschitz boundary $\partial\Omega = \Gamma_1 \cup \Gamma_2$, where Γ_1 is the impervious part of the boundary, Γ_2 is the part in contact with either air or the fluid reservoirs.

The fluid infiltrates through Ω obeying to Darcy's law

$$v = -a(x).\nabla(u + x_n),$$

where $a(x) = (a_{ij}(x))_{ij}$ is the $n \times n$ permeability matrix of the medium, $x = (x_1, ..., x_n)$, v is the fluid velocity and u its pressure.

We are concerned with the problem of finding the pressure u and the saturation χ of the fluid inside Ω. Using the mass conservation law, Darcy's law, the boundary conditions and the initial data, we obtain the following strong formulation for our problem (see [3]):

$$(\mathbf{SF}) \quad \begin{cases} u \geq 0,\ 0 \leq \chi \leq 1,\ u(1-\chi) = 0 & \text{in } Q \\ div(a(x)(\nabla u + \chi e)) - (\alpha u + \chi)_t = 0 & \text{in } Q \\ u = \phi & \text{on } \Sigma_2 \\ (\alpha u + \chi)(\cdot, 0) = \alpha u_0 + \chi_0 & \text{in } \Omega \\ a(x)(\nabla u + \chi e) \cdot \nu = 0 & \text{on } \Sigma_1 \\ a(x)(\nabla u + \chi e) \cdot \nu \leq 0 & \text{on } \Sigma_4 \end{cases}$$

where α, T are positive numbers, $Q = \Omega \times (0,T)$, $\Sigma_1 = \Gamma_1 \times (0,T)$ is the impervious part of $\partial\Omega$, $\Sigma_2 = \Gamma_2 \times (0,T)$ is the pervious part, $\Sigma_3 = \Sigma_2 \cap \{\phi > 0\}$ is the part covered by fluid, and $\Sigma_4 = \Sigma_4 \cap \{\phi = 0\}$ is the part where the fluid flows outside Ω. ϕ is a nonnegative Lipschitz continuous function defined in $\overline{Q}$, ν is the outward unit normal vector to $\partial\Omega$, $e = (0, ..., 0, 1) \in \mathbb{R}^n$, $u_0, \chi_0 : \Omega \longrightarrow \mathbb{R}$ are functions satisfying for a positive constant M

$$0 \leq u_0(x) \leq M, \quad 0 \leq \chi_0(x) \leq 1 \quad \text{for a.e. } x \in \Omega. \tag{1.1}$$

For $a(x)$, we assume that we have for two positive constants λ and Λ

$$\forall \xi \in \mathbb{R}^n, \quad \text{for a.e. } x \in \Omega \quad \lambda|\xi|^2 \leq a(x).\xi.\xi. \tag{1.2}$$

$$\forall \xi \in \mathbb{R}^n, \quad \text{for a.e. } x \in \Omega \quad |a(x).\xi| \leq \Lambda|\xi|. \tag{1.3}$$

Moreover, we assume that

$$div(a(x)e) \in L^2(\Omega). \tag{1.4}$$

Using the strong formulation, we are led to the following weak formulation

$$(\mathbf{P}) \quad \begin{cases} \quad \text{Find } (u,\chi) \in L^2(0,T;H^1(\Omega)) \times L^\infty(Q) \text{ such that :} \\ (i) \quad u \geq 0,\ 0 \leq \chi \leq 1,\ u.(1-\chi) = 0 \quad \text{a.e. in } Q \\ (ii) \quad u = \phi \quad \text{on } \Sigma_2 \\ (iii) \quad \displaystyle\int_Q \big[a(x)\big(\nabla u + \chi e\big)\cdot\nabla\xi - (\alpha u + \chi)\xi_t\big]\,dx\,dt \\ \qquad\qquad \leq \displaystyle\int_\Omega (\chi_0 + \alpha u_0)\xi(x,0)\,dx \\ \quad \forall \xi \in H^1(Q),\ \xi = 0 \text{ on } \Sigma_3,\ \xi \geq 0 \text{ on } \Sigma_2, \\ \quad \xi(x,T) = 0 \text{ for a.e. } x \in \Omega. \end{cases}$$

For the existence of a solution of the problem (P) in the homogeneous case ($a(x) = I_n$), we refer to [2] and [1] respectively in the incompressible ($\alpha = 0$) and compressible ($\alpha > 0$) cases. For the heterogeneous case, we refer to [16] in a more general framework under assumptions (1.1)-(1.4) for both incompressible and compressible cases. For the incompressible case with nonlinear Darcy's law, we refer to [11], [12] and [3] respectively for Dirichlet, Neuman and generalized boundary conditions. Regarding regularity of the solution, we refer to [2] and [1], where it has been proved when $a(x) = I_n$ that $\chi \in C^0([0,T];L^p(\Omega))$ for all $p \geq 1$ in both incompressible and compressible cases, and that $u \in C^0([0,T];L^p(\Omega))$ for all $1 \leq p \leq 2$ in the compressible case. Extensions to the quasilinear and incompressible case were obtained in [11], [12] and [13] in both homogeneous and nonhomogeneous frameworks. The authors of this paper were recently able to extend the above regularity result in [14] to a more general framework under weaker assumptions on the data.

In this paper, we are mainly concerned with the uniqueness of the solution of the problem (P). This question was first addressed for a rectangular homogeneous dam in [15] and [7] respectively for a formulation based on quasi-variational inequalities and for the formulation (P) with a dam wet at the bottom and dry near to the top in the second case. Uniqueness of the solution for a homogeneous dam with general geometry was established by the method of doubling variables in [1], but it is not obvious wether it works in the heterogeneous situation. Extensions to a quasilinear operator modeling incompressible fluid flow governed by a nonlinear Darcy's law with Dirichlet, or Neuman boundary conditions were obtained in [11] and [12] respectively.

Our main result in this work is the uniqueness of problem (P) solution for a heterogeneous and rectangular porous medium assuming it remains wet near the bottom and dry close to the top. Our method is inspired by an idea from [7] in the homogeneous case and relies on solution regularity that has been recently obtained in [14]. Our uniqueness result is new in the heterogeneous and rectangular framework, but most likely the technic is limited to that particular geometry like shape.

2 Preliminary Results

In this work we shall be interested in the following situation of a two dimensional rectangular dam $\Omega = (0,L) \times (0,K)$, with $L, K > 0$, and $\Gamma_1 = [0,L] \times \{0\}$, $\Gamma_2 = (\{0\} \times [0,K]) \cup ([0,L] \times \{K\}) \cup (\{L\} \times [0,K])$ (see Figure 1).
We also assume that

$$a(x)e \in C^{0,1}(\overline{\Omega}), \tag{2.1}$$

$$div(a(x)e) \geq 0 \quad \text{a.e. in } \Omega, \tag{2.2}$$

$$\phi_0 \leq \phi \leq \phi_1 \quad \text{on } \Sigma_2, \tag{2.3}$$

where ϕ_0 and ϕ_1 are two nonnegative Lipschitz continuous functions defined on $\overline{\Omega}$



Figure 1

and satisfying for some $\epsilon_0 > 0$ small enough

$$\begin{cases} \phi_0(0,x_2) = \phi_0(L,x_2) = (\epsilon_0 - x_2)^+ \\ \phi_1(0,x_2) = \phi_1(L,x_2) = (K - \epsilon_0 - x_2)^+ \\ \phi_0(x_1,K) = \phi_1(x_1,K) = 0. \end{cases} \tag{2.4}$$

Let us now denote by (v_i, γ_i) the solution of the stationary problem corresponding to $\phi_i, \quad i = 0, 1$ (see [5])

$$(\mathbf{P_i^s}) \begin{cases} \text{Find } (v_i, \gamma_i) \in H^1(\Omega) \times L^\infty(\Omega) \text{ such that :} \\ \quad (i) \quad v_i \geq 0,\ 0 \leq \gamma_i \leq 1,\ v_i.(1-\gamma_i) = 0 \quad \text{ a.e. in } \Omega \\ \quad (ii) \quad v_i = \phi_i \quad \text{ on } \Gamma_2 \\ \quad (iii) \displaystyle\int_\Omega a(x)(\nabla v_i + \gamma_i e).\nabla \xi dx \leq 0 \\ \qquad \forall \xi \in H^1(\Omega), \quad \xi = 0 \ \text{ on } \ \Gamma_2 \cap \{\phi_i > 0\}, \quad \xi \geq 0 \ \text{ on } \ \Gamma_2 \cap \{\phi_i = 0\}. \end{cases}$$

We recall the following uniqueness result.

Theorem 2.1. *Assume that (2.1)-(2.2) hold. Then the solution (v_i, γ_i) of (P_i^s) is unique and satisfies*

$$\gamma_i = \chi_{\{v_i > 0\}}. \tag{2.5}$$

Proof. When $a(x)e \in C^1(\overline{\Omega})$, we refer to [4], since we cannot have pools in a rectangular dam. One can also argue as in [6]. When $a(x)e \in C^{0,1}(\overline{\Omega})$, one may combine Theorem 5.1 of [5] and the proof of Theorem 6.3 in [4] to establish the result.

□

The following properties for the solution of (P_i^s) $(i = 0, 1)$ hold.

Theorem 2.2. *Assume that (2.1)-(2.4) hold. Then we have*

$$(\epsilon_0 - x_2)^+ \leq v_0 \leq v_1 \leq (K - \epsilon_0 - x_2)^+ \quad a.e. \ in \ \ \Omega \tag{2.6}$$

$$\gamma_0 = 1 \quad a.e. \ in \ \ \Omega \cap \{0 < x_2 < \epsilon_0\} \tag{2.7}$$

$$\gamma_1 = 0 \quad a.e. \ in \ \ \Omega \cap \{K - \epsilon_0 < x_2 < K\}. \tag{2.8}$$

Proof. First, we remark that for each $k \in (0, K)$, $(k - x_2)^+$ satisfies the equation

$$\int_\Omega a(x)(\nabla(k - x_2)^+ + \chi_{\{(k-x_2)^+>0\}}e).\nabla\xi dx = \int_{\Omega\cap\{x_2<k\}} a(x)(-e + e).\nabla\xi dx = 0.$$

It follows that $((k - x_2)^+, \chi_{\{(k-x_2)^+>0\}})$ is a solution of the stationary dam problem for the boundary Dirichlet data $(k - x_2)^+$ on Γ_2. Adapting the proof of uniqueness in [4] and arguing as in [10], and using (2.1)-(2.4), we obtain

$$(\epsilon_0 - x_2)^+ \leq v_0 \leq v_1 \leq (K - \epsilon_0 - x_2)^+ \quad \text{in} \ \ \Omega$$

$$\chi_{\{(\epsilon_0-x_2)^+>0\}} \leq \gamma_0 \leq \gamma_1 \leq \chi_{\{(K-\epsilon_0-x_2)^+>0\}} \quad \text{a.e. in} \ \ \Omega.$$

Hence (2.6)-(2.8) follow. □

Remark 2.1. *Theorems 2.1 and 2.2 remain true without the regularity assumption (2.1) provided the following assumptions on the permeability matrix hold (see [10])*

$$a_{12} = 0 \quad a.e. \ in \ \Omega,$$

$$\frac{\partial a_{22}}{\partial x_2} \geq 0 \quad in \ \mathcal{D}'(\Omega).$$

Next we will construct a solution corresponding to a dam that is wet up to $x_2 = \epsilon_0$ and dry above $x_2 = K - \epsilon_0$ over the whole interval $[0, T]$.

Lemma 2.1. *Assume that (2.1)-(2.4) hold and the initial data satisfies*

$$v_0 \leq u_0 \leq v_1 \quad a.e.\ \ in\ \Omega. \tag{2.9}$$

$$\gamma_0 \leq \chi_0 \leq \gamma_1 \quad a.e.\ \ in\ \Omega. \tag{2.10}$$

Then there exists a solution (u, χ) of problem (P) such that

$$v_0 \leq u \leq v_1 \quad a.e.\ \ in\ Q \tag{2.11}$$

$$\gamma_0 \leq \chi \leq \gamma_1 \quad a.e.\ \ in\ Q. \tag{2.12}$$

Proof. Let $v_{i\epsilon}$ be the solution of the approximating problem of the stationary problem $(P_i^s),\ i = 0, 1$

$$(\mathbf{P^s_{i\epsilon}}) \begin{cases} \text{Find } v_{i\epsilon} \in H^1(\Omega) \text{ such that :} \\ (i) \quad v_{i\epsilon} = \phi_i \quad \text{ on } \Gamma_2 \\ (ii) \quad \displaystyle\int_\Omega a(x)(\nabla v_{i\epsilon} + H_\epsilon(v_{i\epsilon})e).\nabla \xi dx = 0 \\ \quad \forall \xi \in H^1(\Omega), \ \ \xi = 0 \quad \text{ on } \Gamma_2, \end{cases}$$

where $H_\epsilon(s) = \min(1, s^+/\epsilon)$ is an approximation of the Heaviside graph $H(s) = [0, 1]\chi_{\{0\}} + \chi_{(0,\infty)}$.

Let u_ϵ be the solution of the following approximating problem of the problem (P)

$$(\mathbf{P}_\epsilon) \begin{cases} \text{Find } u_\epsilon \in H^1(Q) \text{ such that :} \\ (i) \quad u_\epsilon = \phi \text{ on } \Sigma_2 \\ (ii) \quad \displaystyle\int_Q \big[a(x)\big(\nabla u_\epsilon + H_\epsilon(u_\epsilon)e\big) \cdot \nabla \xi + \epsilon u_{\epsilon t}\xi_t - G_\epsilon(u_\epsilon)\xi_t\big] dxdt \\ \qquad + \displaystyle\int_\Omega G_\epsilon(u_\epsilon(x, T))\xi(x, T)dx = \int_\Omega (\alpha u_{0\epsilon}(x) + \chi_{0\epsilon}(x))\xi(x, 0)dxdy \\ \quad \forall \xi \in H^1(Q), \ \xi = 0 \text{ on } \Sigma_2, \end{cases}$$

where $u_{0\epsilon} = \min(u_0, v_{1\epsilon})$ and $\chi_{0\epsilon} = \min(\chi_0, H_\epsilon(v_{1\epsilon}))$.

If $\xi \in H^1(Q),\ \xi = 0$ on Σ_2, we have from $(P^s_{1\epsilon})(ii)$:

$$\begin{aligned}&\int_Q \big[a(x)\big(\nabla v_{1\epsilon} + H_\epsilon(v_{1\epsilon})e\big)\cdot\nabla\xi - (\alpha v_{1\epsilon} + H_\epsilon(v_{1\epsilon}))\xi_t\big]dxdt\\&+\int_\Omega(\alpha v_{1\epsilon} + H_\epsilon(v_{1\epsilon}))\xi(x,T)dx = \int_\Omega(\alpha v_{1\epsilon} + H_\epsilon(v_{1\epsilon}))\xi(x,0)dx.\end{aligned} \tag{2.13}$$

For $\delta > 0$, the function $\xi_\delta = \dfrac{(u_\epsilon - v_{1\epsilon} - \delta)^+}{u_\epsilon - v_{1\epsilon}}$ belongs to $H^1(Q)$ and satisfies $\xi_\delta = 0$ on Σ_2 since $\phi \le \phi_1$ on Σ_2. Writing (2.13) and $(P_\epsilon)(ii)$ for $\xi = \xi_\delta$ and subtracting the two identities from each other, we get by taking into account (2.9)-(2.10)

$$\begin{aligned}&\int_Q \Big[a(x)\big(\nabla(u_\epsilon - v_{1\epsilon}) + (H_\epsilon(u_\epsilon) - H_\epsilon(v_{1\epsilon}))e\big).\nabla\xi_\delta + \epsilon(u_\epsilon - v_{1\epsilon})_t\xi_{\delta t}\\&-(\alpha(u_\epsilon - v_{1\epsilon}) + H_\epsilon(u_\epsilon) - H_\epsilon(v_{1\epsilon}))\xi_{\delta t}\Big]dxdydt\\&+\int_\Omega(\alpha(u_\epsilon(x,T) - v_{1\epsilon}) + H_\epsilon(u_\epsilon(x,T)) - H_\epsilon(v_{1\epsilon}))\xi_\delta(x,T)dx\\&=\int_\Omega(\alpha(u_{0\epsilon} - v_{1\epsilon}) + \chi_{0\epsilon} - H_\epsilon(v_{1\epsilon}))\xi_\delta(x,0)dx \le 0.\end{aligned} \tag{2.14}$$

By Lemma 2.1 of [16], we obtain from (2.14)

$$u_\epsilon \le v_{1\epsilon} \quad \text{a.e. in } Q \tag{2.15}$$

and by the monotonicity of H_ϵ, we get

$$H_\epsilon(u_\epsilon) \le H_\epsilon(v_{1\epsilon}) \quad \text{a.e. in } Q. \tag{2.16}$$

We recall that from the proof of existence (see [8] or [16] for example), we know that we have up to a subsequence

$$u_\epsilon \rightharpoonup u \qquad \text{weakly in } L^2(0,T;H^1(\Omega)). \tag{2.17}$$

$$H_\epsilon(u_\epsilon) \rightharpoonup \chi \qquad \text{weakly in } L^2(Q) \tag{2.18}$$

where (u,χ) is a solution of problem (P).

Similarly, we have since the solution of problem (P_1^s) is unique

$$v_{1\epsilon} \rightharpoonup v_1 \qquad \text{weakly in } H^1(\Omega). \tag{2.19}$$

$$H_\epsilon(v_{1\epsilon}) \rightharpoonup \gamma_1 \qquad \text{weakly in } L^2(\Omega). \tag{2.20}$$

Now, let $\xi \in \mathcal{D}(Q)$ with $\xi \geq 0$. Passing to the limit, we obtain by using (2.15)-(2.20)

$$\begin{aligned} \int_Q (v_1 - u)\xi dxdt &= \lim_{\epsilon\to 0} \int_Q (v_{1\epsilon} - u_\epsilon)\xi dxdt \geq 0, \\ \int_Q (\gamma_1 - \chi)\xi dxdt &= \lim_{\epsilon\to 0} \int_Q (H_\epsilon(v_{1\epsilon}) - H_\epsilon(u_\epsilon))\xi dxdt \geq 0, \end{aligned}$$

which leads to

$$u \leq v_1 \quad \text{a.e. in } Q \tag{2.21}$$

$$\chi \leq \gamma_1 \quad \text{a.e. in } Q. \tag{2.22}$$

Similarly, for $\delta > 0$ the function $\xi_\delta = \dfrac{(v_{0\epsilon} - u_\epsilon - \delta)^+}{v_{0\epsilon} - u_\epsilon}$ belongs to $H^1(Q)$ and satisfies $\xi_\delta = 0$ on Σ_2 since $\phi_0 \leq \phi$ on Σ_2. Then by taking into account (2.9)-(2.10), we get

$$\begin{aligned} &\int_Q \Big[a(x)\big(\nabla(v_{0\epsilon} - u_\epsilon) + (H_\epsilon(v_{0\epsilon}) - H_\epsilon(u_\epsilon))e\big).\nabla\xi_\delta + \epsilon(v_{0\epsilon} - u_\epsilon)_t\xi_{\delta t} \\ &-(\alpha(v_{0\epsilon} - u_\epsilon) + H_\epsilon(v_{0\epsilon}) - H_\epsilon(u_\epsilon))\xi_{\delta t}\Big] dxdt \\ &+ \int_\Omega (\alpha(v_{0\epsilon} - u_\epsilon(x,T)) + H_\epsilon(v_{0\epsilon}) - H_\epsilon(u_\epsilon(x,T)))\xi_\delta(x,T)dx \\ &= \int_\Omega (\alpha(v_{0\epsilon} - u_{0\epsilon}) + H_\epsilon(v_{0\epsilon}) - \chi_{0\epsilon})\xi_\delta(x,0)dx \leq 0. \end{aligned} \tag{2.23}$$

By Lemma 2.1 of [16], we obtain from (2.23)

$$v_{0\epsilon} \leq u_\epsilon \quad \text{a.e. in } Q \tag{2.24}$$

and by the monotonicity of H_ϵ, we get

$$H_\epsilon(v_{0\epsilon}) \leq H_\epsilon(u_\epsilon) \quad \text{a.e. in } Q. \tag{2.25}$$

Arguing as above and using (2.24)-(2.25), we obtain by passing to the limit up to a subsequence, that we have for any $\xi \in \mathcal{D}(Q)$ with $\xi \geq 0$

$$\int_Q (u - v_0)\xi dxdt = \lim_{\epsilon \to 0} \int_Q (u_\epsilon - v_{0\epsilon})\xi dxdt \geq 0,$$
$$\int_Q (\chi - \gamma_0)\xi dxdt = \lim_{\epsilon \to 0} \int_Q (H_\epsilon(u_\epsilon) - H_\epsilon(v_{0\epsilon}))\xi dxdt \geq 0,$$

which leads to

$$v_0 \leq u \quad \text{a.e. in } Q \tag{2.26}$$

$$\gamma_0 \leq \chi \quad \text{a.e. in } Q. \tag{2.27}$$

Combining (2.21)-(2.22) and (2.26)-(2.27), we obtain (2.11)-(2.12). □

Remark 2.2. *Assume that $a(x)e \in C^{0,1}(\overline{\Omega})$. Then we get from (2.11)-(2.12) taking into account (2.5)*

$$u(x,t) > 0 \quad \textit{if} \;\; 0 < x_2 < \epsilon_0 \tag{2.28}$$

$$u(x,t) = \chi(x,t) = 0 \quad \textit{if} \;\; K - \epsilon_0 < x_2 < K. \tag{2.29}$$

3 Uniqueness of the solution in rectangular dams

In this section we assume that

$$a(x) \in C^{0,1}(\overline{\Omega}), \quad \text{with} \quad N = \sup_{i,j,k} |(a_{ij})_{x_k}|_\infty. \tag{3.1}$$

$$a(x) \text{ is a symmetric matrix.} \tag{3.2}$$

Here is our main result.

Theorem 3.1. *Assume that (2.2) and (3.1)-(3.2) hold. Then the solution of the problem (P) associated with the initial data (u_0, χ_0) and satisfying (2.28)-(2.29) is unique.*

Let (u_1, χ_1) and (u_2, χ_2) be two solutions of the problem (P) satisfying (2.28)-(2.29). Set

$$w = u_1 - u_2 \quad \text{and} \quad \eta = \alpha w + \chi_1 - \chi_2.$$

We consider the following problem

$$\begin{aligned}
&\text{Find } v \in L^2(0,T;H^1(\Omega)) \ \text{ such that :} \\
&div(a(x)\nabla v) = -\eta \ \text{ in } \ \Omega \ \text{ for each } t \in [0,T] && (3.3)\\
&v = 0 \ \text{ on } \ \Gamma_2 && (3.4)\\
&a(x)\nabla v.\nu = 0 \ \text{ on } \ \Gamma_1. && (3.5)
\end{aligned}$$

The we have

Lemma 3.1. *There exists a unique weak solution of the problem (3.3)-(3.5).*

Proof. First, we observe (see [14]) that $\alpha u_i + \chi_i \in C^0([0,T];L^2(\Omega))$, $i = 0, 1$. As a consequence, we have $\eta \in C^0([0,T];L^2(\Omega))$. Let $V = \{v \in H^1(\Omega) \ / \ v = 0 \text{ on } \Gamma_2 \ \}$. Then V is a Hilbert space under the $H^1(\Omega)$ norm, and by applying Lax-Milgram's Theorem and taking into account (1.2)-(1.3), there exists for each $t \in [0,T]$ a unique solution $v(x,t)$ of the following problem

$$\begin{cases} v(.,t) \in V \\ \displaystyle\int_\Omega a(x)\nabla v(x,t).\nabla \xi dx = \int_\Omega \eta(x,t)\xi dx \quad \forall \xi \in V. \end{cases} \tag{3.6}$$

Choosing $\xi \in \mathcal{D}(\Omega)$ in (3.6), we obtain (3.3) in $\mathcal{D}'(\Omega)$ and therefore in $C^0([0,T];L^2(\Omega))$. (3.4) is satisfied in the trace sense and (3.5). Writing (3.6) for $\xi \in C^\infty(\overline{\Omega})$ with $\xi = 0$ on Γ_2, and taking into account (3.3), we obtain (3.4) in $H^{-1/2}(\Gamma_1)$.

Choosing v as a test function in (3.6) and using (1.2), Hölder and Poincaré's inequalities, we obtain

$$\int_\Omega |\nabla v(x,t)|^2 dx \leq \frac{1}{\lambda^2} \int_\Omega |\eta(x,t)|^2 dx. \tag{3.7}$$

Integrating (3.7) over the interval $[0, T]$ and using the fact that $\chi_i \in L^\infty(Q)$, $u_i \in L^\infty(0, T; L^\infty(\Omega))$ (see [14]), we obtain

$$\int_Q |\nabla v(x,t)|^2 dxdt \leq \frac{1}{\lambda^2} \int_0^T \int_\Omega |\eta(x,t)|^2 dxdt \leq \frac{|\eta|_{L^\infty(0,T;L^\infty(\Omega))}}{\lambda^2}.$$

Using Poincaré's inequality, we obtain $v \in L^2(0, T; H^1(\Omega))$. Hence v is the unique solution of (3.3)-(3.5).

$\square$

Remark 3.1. *By the regularity theory (see [9] for example), the solution v of the problem (3.3)-(3.5) satisfies $v \in L^2(0, T; C^1(\Omega \cup \mathring{\Gamma}_1 \cup \mathring{\Gamma}_2)) \cap L^2(0, T; H^2(\Omega \cup \mathring{\Gamma}_1 \cup \mathring{\Gamma}_2))$.*

Now, let us denote by $\widetilde{g}$ the mean with respect to t of a function $g(x, t)$ defined by

$$\widetilde{g}(x,t) = \frac{1}{h} \int_t^{t+h} g(x,s) ds.$$

Then we have

$$\begin{aligned} &\widetilde{g} \to g \quad \text{as} \quad h \to 0 \\ &\frac{\partial \widetilde{g}}{\partial t} = \frac{1}{h}(g(x,t+h) - g(x,t)). \end{aligned} \tag{3.8}$$

Moreover, it is easy to check that

$$div(a(x)\nabla \widetilde{v}) = -\widetilde{\eta} \quad \text{in} \quad \Omega \quad \text{for all} \quad t \in [0, T] \tag{3.9}$$

$$\widetilde{v} = 0 \quad \text{on} \quad \Gamma_2 \tag{3.10}$$

$$a(x)\nabla \widetilde{v}.\nu = 0 \quad \text{on} \quad \Gamma_1. \tag{3.11}$$

Since $\chi_i = 1$ $(i = 1, 2)$ in a neighborhood of Γ_1, we obtain from $(P)iii)$

$$\begin{aligned} &(\alpha u_i + \chi_i)_t = div(a(x)(\nabla u_i + \chi_i e)) \quad \text{in } \mathcal{D}'(Q), \\ &a(x)\big(\nabla u_i + e\big).\nu = 0 \quad \text{on} \quad \Gamma_1, \quad i = 1, 2. \end{aligned}$$

Using the fact that $u_1 = u_2$ on Σ_2, and writing the previous two equations for (u_1, χ_1) and (u_2, χ_2) and subtracting them from each other, we get

$$(\alpha w + \chi_1 - \chi_2)_t = div\big(a(x)(\nabla w + (\chi_1 - \chi_2)e)\big) \quad \text{in } \mathcal{D}'(Q). \tag{3.12}$$

$$w = 0 \ \text{ on } \ \Sigma_2 \tag{3.13}$$

$$a(x)(\nabla w).\nu = 0 \ \text{ on } \ \Sigma_1. \tag{3.14}$$

Then we have

Lemma 3.2. *For h small enough we have*

$$\widetilde{\eta}_t = div(a(x)(\nabla \widetilde{w} + (\widetilde{\chi_1} - \widetilde{\chi_2})e)) \quad \textit{in } \mathcal{D}'(Q) \tag{3.15}$$

$$\widetilde{w} = 0 \ \textit{ on } \ \Sigma_2 \tag{3.16}$$

$$a(x)\nabla \widetilde{w}.\nu = 0 \ \textit{ on } \ \Sigma_1. \tag{3.17}$$

Proof. (3.16) and (3.17) are a direct consequence of (3.13) and (3.14). To establish (3.15), let $\zeta \in \mathcal{D}(Q)$ such that for some $\tau_0 > 0$, $supp(\zeta) \subset \Omega \times (\tau_0, T - \tau_0)$. We denote by $\widehat{\zeta}$ the function defined by $\widehat{\zeta}(x,t) = \frac{1}{h}\int_{t-h}^{t} \zeta(x,s)ds$. Since for $|h| < \tau_0/2$, the functions $\pm\widehat{\zeta}$ are test functions for problem (P), we obtain for $i = 1, 2$

$$\int_Q a(x)\big(\nabla u_i + \chi_i e\big) \cdot \nabla\widehat{\zeta} dxdt = \int_Q (\alpha u_i + \chi_i)\widehat{\zeta}_t dxdt. \tag{3.18}$$

For the right hand side of (3.18), we have by using change of variables

$$\begin{aligned}
\int_Q (\alpha u_i + \chi_i)\widehat{\zeta}_t dxdt &= \int_Q (\alpha u_i + \chi_i)\frac{1}{h}\zeta(x,t)dxdt - \int_Q (\alpha u_i + \chi_i)\frac{1}{h}\zeta(x,t-h)dxdt \\
&= \int_Q (\alpha u_i + \chi_i)(x,t)\frac{1}{h}\zeta(x,t)dxdt - \int_\Omega \int_{-h}^{T-h} \frac{1}{h}(\alpha u_i(x,t+h) + \chi_i(x,t+h))\zeta(x,t)dxdt \\
&= \int_Q (\alpha u_i + \chi_i)(x,t)\frac{1}{h}\zeta(x,t)dxdt - \int_\Omega \int_{0}^{T} \frac{1}{h}(\alpha u_i(x,t+h) + \chi_i(x,t+h))\zeta(x,t)dxdt \\
&= -\int_Q \frac{1}{h}[(\alpha u_i(x,t+h) + \chi_i(x,t+h)) - (\alpha u_i + \chi_i)(x,t)]\zeta(x,t)dxdt \\
&= -\int_Q (\widetilde{\alpha u_i + \chi_i})_t \zeta(x,t)dxdt.
\end{aligned} \tag{3.19}$$

For the left hand side of (3.18), we have by integrating by parts

$$
\begin{aligned}
\int_Q a(x)\big(\nabla u_i + \chi_i e\big) \cdot \nabla\widehat{\zeta} dxdt &= -\int_Q \Big[\int_0^t a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\widehat{\zeta}_t dxdt \\
&= -\int_Q \Big[\int_0^t a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\big(\frac{1}{h}(\zeta(x,t) - \zeta(x,t-h))\big)dxdt \\
&= -\int_Q \frac{1}{h}\Big[\int_0^t a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\zeta(x,t)dxdt \\
&\quad + \int_Q \frac{1}{h}\Big[\int_0^t a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\zeta(x,t-h)dxdt \\
&= -\int_Q \frac{1}{h}\Big[\int_0^t a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\zeta(x,t)dxdt \\
&\quad + \int_\Omega\int_0^T \frac{1}{h}\Big[\int_0^{t+h} a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\zeta(x,t)dxdt \\
&= \int_Q \frac{1}{h}\Big[\int_t^{t+h} a(x)\big(\nabla u_i + \chi_i e\big)ds\Big] \cdot \nabla\zeta(x,t)dxdt \\
&= \int_Q a(x)\big(\nabla\widetilde{u_i} + \widetilde{\chi_i} e\big) \cdot \nabla\zeta(x,t)dxdt. \qquad (3.20)
\end{aligned}
$$

Then we deduce from (3.18)-(3.20) that

$$\int_Q a(x)\big(\nabla\widetilde{u_i} + \widetilde{\chi_i} e\big) \cdot \nabla\zeta(x,t)dxdt = -\int_Q (\alpha\widetilde{u_i + \chi_i})_t\zeta(x,t)dxdt.$$

Writing the last equation for $i = 1, 2$ and subtracting the two equations, we get

$$\int_Q a(x)\big(\nabla\widetilde{w} + (\widetilde{\chi_1} - \widetilde{\chi_2})e\big) \cdot \nabla\zeta(x,t)dxdt = -\int_Q \widetilde{\eta}_t\zeta(x,t)dxdt$$

which is (3.15). □

To prove Theorem 3.1, we need two more lemmas.

Lemma 3.3.

$$
\begin{aligned}
&\frac{1}{2}\frac{\partial}{\partial t}\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx + \int_\Omega \widetilde{w}\widetilde{\eta}dx = \int_\Omega \alpha\widetilde{w}(a_{11}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})dx \\
&+ \int_\Omega (a_{11}\widetilde{v}_{x_1} + a_{12}\widetilde{v}_{x_2})_{x_1}(a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})dx + \int_\Omega (a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})_{x_2}(a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})dx. \qquad (3.21)
\end{aligned}
$$

Proof. From (3.9) and (3.15) we derive

$$-div(a(x)\nabla\widetilde{v})_t = div(a(x)(\nabla\widetilde{w} + (\widetilde{\chi_1} - \widetilde{\chi_2})e)) \quad \text{in } \mathcal{D}'(\Omega). \tag{3.22}$$

Using (3.10)-(3.11), and taking into account (3.2), we obtain

$$\begin{aligned}< -div(a(x)(\nabla\widetilde{v})_t, \widetilde{v} >&= \int_\Omega a(x)\nabla\widetilde{v}_t.\nabla\widetilde{v}dx - \int_{\partial\Omega} a(x)\nabla\widetilde{v}_t.\nu\ \widetilde{v}d\sigma(x)\\ &= \int_\Omega a(x)\nabla\widetilde{v}_t.\nabla\widetilde{v}dx = \frac{1}{2}\frac{\partial}{\partial t}\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx. \end{aligned}\tag{3.23}$$

Similarly, we get by using (3.10), (3.17), and taking into account (3.2)

$$\begin{aligned}< div(a(x)\nabla\widetilde{w}), \widetilde{v} > &= -\int_\Omega a(x)\nabla\widetilde{w}.\nabla\widetilde{v}dx + \int_{\partial\Omega} a(x)\nabla\widetilde{w}.\nu\ \widetilde{v}d\sigma(x)\\ &= -\int_\Omega a(x)\nabla\widetilde{w}.\nabla\widetilde{v}dx = -\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{w}dx\end{aligned}$$

which can be written using (3.9) as

$$< div(a(x)\nabla\widetilde{w}), \widetilde{v} > \ = \ -\int_\Omega \widetilde{w}\widetilde{\eta}dx. \tag{3.24}$$

Similarly, using (3.10) and the fact that u_1, u_2 satisfy (3.28)-(3.29), we get for $\Omega_0 = (0, L) \times (\epsilon_0, K - \epsilon_0)$

$$\begin{aligned}&< div((\widetilde{\chi_1} - \widetilde{\chi_2})a(x)e), \widetilde{v} >= -\int_{\Omega_0} (\widetilde{\chi_1} - \widetilde{\chi_2})a(x)e.\nabla\widetilde{v}dx\\ &= \int_{\Omega_0} (-\widetilde{\eta} + \alpha\widetilde{w})a(x)e.\nabla\widetilde{v}dx\\ &= \int_{\Omega_0} (div(a(x)\nabla\widetilde{v}) + \alpha\widetilde{w})a(x)e.\nabla\widetilde{v}dx\\ &= \int_{\Omega_0} div(a(x)\nabla\widetilde{v}))a(x)e.\nabla\widetilde{v}dx + \int_{\Omega_0} \alpha\widetilde{w}a(x)e.\nabla\widetilde{v}dx\\ &= \int_{\Omega_0} (a_{11}\widetilde{v}_{x_1} + a_{12}\widetilde{v}_{x_2})_{x_1}(a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})dx\\ &+ \int_{\Omega_0} (a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})_{x_2}(a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})dx\\ &+ \int_{\Omega_0} \alpha\widetilde{w}(a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})dx.\end{aligned}\tag{3.25}$$

Hence by combining (3.22)-(3.25), we get (3.21). □

Lemma 3.4. *There exists a positive constant C such that*

$$\frac{\partial}{\partial t}\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx + 2\int_\Omega \widetilde{w}(\widetilde{\chi_1}-\widetilde{\chi_2})dx \leq C\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx. \tag{3.26}$$

The proof of Lemma 3.4 requires a lemma.

Lemma 3.5. *There exists a positive constant C such that*

$$\int_{\epsilon_0}^{K-\epsilon_0}|\nabla\widetilde{v}(0,x_2)|^2dx_2 \leq C\int_\Omega|\nabla\widetilde{v}|^2dx \tag{3.27}$$

$$\int_{\epsilon_0}^{K-\epsilon_0}|\nabla\widetilde{v}(L,x_2)|^2dx_2 \leq C\int_\Omega|\nabla\widetilde{v}|^2dx \tag{3.28}$$

$$\int_0^L|\nabla\widetilde{v}(x_1,\epsilon_0)|^2dx_1 \leq C\int_\Omega|\nabla\widetilde{v}|^2dx \tag{3.29}$$

$$\int_0^L|\nabla\widetilde{v}(x_1,K-\epsilon_0)|^2dx_1 \leq C\int_\Omega|\nabla\widetilde{v}|^2dx. \tag{3.30}$$

Proof. *i*) Since $\widetilde{v}\in C^1(\overline{\Omega}_0)$, we have $\int_{\epsilon_0}^{K-\epsilon_0}|\nabla\widetilde{v}(0,x_2)|^2dx_2 = \lim_{\delta\to 0}\frac{1}{\delta}\int_0^\delta\int_{\epsilon_0}^{K-\epsilon_0}|\nabla\widetilde{v}|^2dx.$ It follows that we have for $\delta_1>0$ small enough

$$\int_{\epsilon_0}^{K-\epsilon_0}|\nabla\widetilde{v}(0,x_2)|^2dx_2 \leq \frac{2}{\delta_1}\int_0^{\delta_1}\int_{\epsilon_0}^{K-\epsilon_0}|\nabla\widetilde{v}|^2dx$$

which gives (3.27).

In the same way we establish (3.28), (3.29), and (3.30). □

Proof of Lemma 3.4. We shall estimate the three integrals in the right hand side of (3.21). First, we obtain by applying Young's inequality and using (1.2)-(1.3)

$$\begin{aligned}\int_{\Omega_0}\alpha\widetilde{w}(a_{12}\widetilde{v}_{x_1}+a_{22}\widetilde{v}_{x_2})dx &\leq \frac{\alpha}{2}\int_{\Omega_0}\widetilde{w}^2dx+\frac{\alpha}{2}\int_{\Omega_0}(a_{12}\widetilde{v}_{x_1}+a_{22}\widetilde{v}_{x_2})^2dx\\ &\leq \frac{\alpha}{2}\int_\Omega\widetilde{w}^2dx+\frac{\alpha}{2}\int_\Omega(a_{12}^2+a_{22}^2)|\nabla\widetilde{v}|^2dx\\ &\leq \frac{\alpha}{2}\int_\Omega\widetilde{w}^{\,2}dx+\frac{\alpha\Lambda^2}{\lambda}\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx. \qquad (3.31)\end{aligned}$$

Next, we have

$$\begin{aligned}
&\int_{\Omega_0}(a_{11}\widetilde{v}_{x_1}+a_{12}\widetilde{v}_{x_2})_{x_1}(a_{12}\widetilde{v}_{x_1}+a_{22}\widetilde{v}_{x_2})dx=\int_{\Omega_0}(a_{11}\widetilde{v}_{x_1})_{x_1}(a_{12}\widetilde{v}_{x_1})dx\\
&\quad+\int_{\Omega_0}(a_{11}\widetilde{v}_{x_1})_{x_1}(a_{22}\widetilde{v}_{x_2})dx+\int_{\Omega_0}(a_{12}\widetilde{v}_{x_2})_{x_1}(a_{12}\widetilde{v}_{x_1})dx+\int_{\Omega_0}(a_{12}\widetilde{v}_{x_2})_{x_1}(a_{22}\widetilde{v}_{x_2})dx\\
&\quad=I_1+I_2+I_3+I_4. \qquad (3.32)
\end{aligned}$$

Let us estimate the integrals I_i. Expanding and integrating by parts, and using (1.2)-(1.3), (3.1), and (3.28), we obtain for a positive constant C_1

$$\begin{aligned}
I_1 &= \int_{\Omega_0}(a_{11})_{x_1}a_{12}\widetilde{v}^2_{x_1}dx+\frac{1}{2}\int_{\Omega_0}a_{11}a_{12}(\widetilde{v}^2_{x_1})_{x_1}dx\\
&= \int_{\Omega_0}(a_{11})_{x_1}a_{12}\widetilde{v}^2_{x_1}dx-\frac{1}{2}\int_{\Omega_0}(a_{11}a_{12})_{x_1}\widetilde{v}^2_{x_1}dx\\
&\quad+\frac{1}{2}\int_{\epsilon_0}^{K-\epsilon_0}(a_{11}a_{12}\widetilde{v}^2_{x_1})(L,x_2)dx_2-\frac{1}{2}\int_{\epsilon_0}^{K-\epsilon_0}(a_{11}a_{12}\widetilde{v}^2_{x_1})(0,x_2)dx_2
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2}\int_{\Omega_0}(a_{11})_{x_1}a_{12}\widetilde{v}^2_{x_1}dx-\frac{1}{2}\int_{\Omega_0}a_{11}(a_{12})_{x_1}\widetilde{v}^2_{x_1}dx+\frac{\Lambda^2}{2}\int_{\epsilon_0}^{K-\epsilon_0}\widetilde{v}^2_{x_1}(L,x_2)dx_2\\
&\leq \Lambda N\int_{\Omega_0}\widetilde{v}^2_{x_1}dx+\frac{\Lambda^2}{2}\int_{\epsilon_0}^{K-\epsilon_0}\widetilde{v}^2_{x_1}(L,x_2)dx_2\leq C_1\int_{\Omega}a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx. \qquad (3.33)
\end{aligned}$$

Note that since $\widetilde{v}(0,x_2)=\widetilde{v}(L,x_2)=0$ for $0<x_2<K$, we have

$$\widetilde{v}_{x_2}(0,x_2)=\widetilde{v}_{x_2}(L,x_2)=0 \quad \text{for} \quad 0<x_2<K. \qquad (3.34)$$

Integrating by parts and using (3.34), we obtain

$$\begin{aligned}
I_2 &= -\int_{\Omega_0}a_{11}\widetilde{v}_{x_1}(a_{22}\widetilde{v}_{x_2})_{x_1}dx=-\int_{\Omega_0}a_{11}(a_{22})_{x_1}\widetilde{v}_{x_1}\widetilde{v}_{x_2}dx-\frac{1}{2}\int_{\Omega_0}a_{11}a_{22}(\widetilde{v}^2_{x_1})_{x_2}dx\\
&= -\int_{\Omega_0}a_{11}(a_{22})_{x_1}\widetilde{v}_{x_1}\widetilde{v}_{x_2}dx+\frac{1}{2}\int_{\Omega_0}a_{11}(a_{22})_{x_2}\widetilde{v}^2_{x_1}dx+\frac{1}{2}\int_{\Omega_0}(a_{11})_{x_2}a_{22}\widetilde{v}^2_{x_1}dx\\
&\quad-\frac{1}{2}\int_0^L(a_{11}a_{22}\widetilde{v}^2_{x_1})(x_1,K-\epsilon_0)dx_1+\frac{1}{2}\int_0^L(a_{11}a_{22}\widetilde{v}^2_{x_1})(x_1,\epsilon_0)dx_1. \qquad (3.35)
\end{aligned}$$

Using Young's inequality, (1.2)-(1.3), (3.1), and (3.29), we obtain from (3.35), for a positive constant C_2

$$\begin{aligned} I_2 &\leq \Lambda N \int_{\Omega_0} |\widetilde{v}_{x_1}|.|\widetilde{v}_{x_2}| dx + \Lambda N \int_{\Omega_0} \widetilde{v}_{x_1}^2 dx + \frac{\Lambda^2}{2} \int_0^L \widetilde{v}_{x_1}^2(x_1, \epsilon_0) dx_1 \\ &\leq \frac{3\Lambda N}{2} \int_{\Omega_0} |\nabla \widetilde{v}|^2 dx + \frac{\Lambda^2}{2} \int_0^L \widetilde{v}_x^2(x_1, \epsilon_0) dx_1 \leq C_2 \int_\Omega a(x) \nabla \widetilde{v} . \nabla \widetilde{v} dx. \end{aligned} \tag{3.36}$$

Expanding and integrating by parts, and using Young's inequality, (1.2)-(1.3), (3.1), and (3.30), we obtain for a positive constant C_3

$$\begin{aligned} I_3 &= \int_{\Omega_0} (a_{12})_{x_1} a_{12} \widetilde{v}_{x_1} \widetilde{v}_{x_2} dx + \frac{1}{2} \int_{\Omega_0} a_{12}^2 (\widetilde{v}_{x_1}^2)_{x_2} dx \\ &= \int_{\Omega_0} (a_{12})_{x_1} a_{12} \widetilde{v}_{x_1} \widetilde{v}_{x_2} dx - \frac{1}{2} \int_{\Omega_0} (a_{12}^2)_{x_2} \widetilde{v}_{x_1}^2 dx \\ &+ \frac{1}{2} \int_0^L (a_{12}^2 \widetilde{v}_{x_1}^2)(x_1, K - \epsilon_0) dx_1 - \frac{1}{2} \int_0^L (a_{12}^2 \widetilde{v}_{x_1}^2)(x_1, \epsilon_0) dx_1 \\ &\leq \frac{\Lambda N}{2} \int_{\Omega_0} (\widetilde{v}_{x_1}^2 + \widetilde{v}_{x_2}^2) dx - \frac{1}{2} \int_{\Omega_0} (a_{12}^2)_{x_2} \widetilde{v}_{x_1}^2 dx + \frac{\Lambda^2}{2} \int_0^L \widetilde{v}_{x_1}^2 (x_1, K - \epsilon_0) dx_1 \\ &\leq C_3 \int_\Omega a(x) \nabla \widetilde{v} . \nabla \widetilde{v} dx. \end{aligned} \tag{3.37}$$

Integrating by parts and using (3.34), and taking into account (1.2)-(1.3) and (3.1), we obtain for a positive constant C_4

$$\begin{aligned} I_4 &= \int_{\Omega_0} (a_{12})_{x_1} a_{22} \widetilde{v}_{x_2}^2 dx + \frac{1}{2} \int_{\Omega_0} a_{12} a_{22} (\widetilde{v}_{x_2}^2)_{x_1} dx \\ &= \int_{\Omega_0} (a_{12})_{x_1} a_{22} \widetilde{v}_{x_2}^2 dx - \frac{1}{2} \int_{\Omega_0} (a_{12} a_{22})_{x_1} \widetilde{v}_{x_2}^2 dx \\ &= \frac{1}{2} \int_{\Omega_0} \big((a_{12})_{x_1} a_{22} - a_{12} (a_{22})_{x_1} \big) \widetilde{v}_{x_2}^2 dx \leq C_4 \int_\Omega a(x) \nabla \widetilde{v} . \nabla \widetilde{v} dx. \end{aligned} \tag{3.38}$$

To estimate the last integral in the right hand side of (3.21), we integrate by parts and use (3.29-(3.30) and (1.2). We obtain for a positive constant C_5

$$\begin{aligned} &\int_{\Omega_0} (a_{12} \widetilde{v}_{x_1} + a_{22} \widetilde{v}_{x_2})_{x_2} (a_{12} \widetilde{v}_{x_1} + a_{22} \widetilde{v}_{x_2}) dx = \frac{1}{2} \int_{\Omega_0} ((a_{12} \widetilde{v}_{x_1} + a_{22} \widetilde{v}_{x_2})^2)_{x_2} dx \\ &= \frac{1}{2} \int_0^L (a_{12} \widetilde{v}_{x_1} + a_{22} \widetilde{v}_{x_2})^2 (x_1, K - \epsilon_0) dx_1 - \frac{1}{2} \int_0^L (a_{12} \widetilde{v}_{x_1} + a_{22} \widetilde{v}_{x_2})^2 (x_1, \epsilon_0) dx \end{aligned}$$

$$\begin{aligned}&\leq \frac{1}{2}\int_0^L (a_{12}\widetilde{v}_{x_1} + a_{22}\widetilde{v}_{x_2})^2(x_1, K-\epsilon_0)dx_1\\ &\leq \int_0^L ((a_{12}\widetilde{v}_{x_1})^2 + (a_{22}\widetilde{v}_{x_2})^2)(x_1, K-\epsilon_0)dx_1\\ &\leq \Lambda^2\int_0^L ((\widetilde{v}_{x_1})^2 + (\widetilde{v}_{x_2})^2)(x_1, K-\epsilon_0)dx_1 \leq C_5\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx. \quad (3.39)\end{aligned}$$

Finally, combining (3.21), (3.31)-(3.33), and (3.35)-(3.39), we get for a positive constant C

$\frac{\partial}{\partial t}\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx + 2\int_\Omega \widetilde{w}(\widetilde{\chi_1} - \widetilde{\chi_2})dx \leq C\int_\Omega a(x)\nabla\widetilde{v}.\nabla\widetilde{v}dx$, which is (3.26). □

Proof of Theorem 3.1. First, integrating (3.26) from 0 to t and letting $h \to 0$, we get

$$\int_\Omega a(x)\nabla v.\nabla vdx + 2\int_0^t\int_\Omega w(\chi_1 - \chi_2)dxds \leq C\int_0^t\int_\Omega a(x)\nabla v.\nabla vdxds. \quad (3.40)$$

Next, we observe that since $u_i \in H(\chi_i)$ a.e. in Q, we have

$$w(\chi_1 - \chi_2) \geq 0 \quad \text{a.e. in} \quad Q. \quad (3.41)$$

Setting $F(t) = \int_0^t\int_\Omega a(x)\nabla v.\nabla vdxds$, we deduce from (3.40)-(3.41) that

$$F'(t) \leq CF(t) \quad \forall t \in [0,T]. \quad (3.42)$$

Integrating (3.42), we get since $F(0) = 0, \quad 0 \leq F(t) \leq F(0)e^{Ct} = 0 \quad \forall t \in [0,T]$, or

$$\int_0^t\int_\Omega a(x)\nabla v.\nabla vdxds = 0 \quad \forall t \in [0,T].$$

Using (1.2), we obtain $\nabla v = 0$ a.e. in Q. Taking into account that $v = 0$ on $\Gamma_2 \subset \partial\Omega$ and the connectedness of Ω, we obtain $v = 0$ in Q. Going back to (3.3), we obtain $\eta = 0$ a.e. in Q, which reads

$$w + \chi_1 - \chi_2 = 0 \quad \text{in} \quad Q. \quad (3.43)$$

Multiplying (3.43) by w, we get $\alpha w^2 + w(\chi_1 - \chi_2) = 0$ a.e. in Q. Taking into account (3.41), we obtain $w^2 = 0$ a.e. in Q, or $u_1 = u_2$ a.e. in Q. Finally, we obtain from (3.43) that $\chi_1 = \chi_2$ a.e. in Q. This achieves the proof. □

Acknowledgments The second author is grateful to Prof. J. F. Rodrigues for kindly inviting him to the CMAF where he enjoyed excellent research conditions during the preparation of his Ph.D Thesis.

References

[1] J. Carrillo, *On the uniqueness of the solution of the evolution dam problem,* Nonlinear Analysis, Theory, Methods and Applications, 22(5), 573-607 (1994).

[2] J. Carrillo and G. Gilardi : La vitesse de propagation dans le problème de la digue. Ann. Fac. Sci. Toulouse Math. (5) 11, No. 3, 7-28 (1990).

[3] J. Carrillo and A. Lyaghfouri : A filtration problem with nonlinear Darcy's law and generalized boundary conditions. *Advances in Differential Equations* Vol. 5, No. 4-6, 515-555 (2000).

[4] Challal, S., Lyaghfouri, A. : A Filtration Problem through a Heterogeneous Porous Medium. Interfaces and Free Boundaries 6, 55-79 (2004).

[5] S. Challal and A. Lyaghfouri : On a class of Free Boundary Problems of type $div(a(X)\nabla u) = -div(H(X)\chi(u))$. Differential and Integral Equations, Vol. 19, No. 5, 481-516 (2006).

[6] S. Challal and A. Lyaghfouri : The Heterogeneous Dam problem with Leaky Boundary Condition. Communications in Pure and Applied Analysis. Vol. 10, No. 1, 93-125 (2011).

[7] E. Dibenedetto and A. Friedam, *Periodic behaviour for the evolutionary dam problem and related free boundary problems,* Communs partial diff. Eqns 11, 1297-1377 (1986).

[8] G. Gilardi : A new approach to evolution free boundary problems, Comm. part. diff. Eqns Vol. 4, 1099-I 123 (1979); Vol. 5, 983-984 (1980).

[9] D. Gilbarg, N.S. Trudinger : Elliptic Partial Differential Equations of Second Order. Springer-Verlag 1983.

[10] A. Lyaghfouri : The Inhomogeneous Dam Problem with Linear Darcy's Law and Dirichlet Boundary Conditions. Mathematical Models and Methods in Applied Sciences 8(6), 1051-1077 (1996).

[11] A. Lyaghfouri : The evolution dam problem with nonlinear Darcy's law and Dirichlet boundary conditions. *Portugaliae Mathematica* 56 (1), 1-38 (1999).

[12] A. Lyaghfouri : *The evolution dam problem with nonlinear Darcy's law and leaky boundary conditions*. Ricerche di Matematica Vol. XLVII, Fasc. 2, 297-357, (1998).

[13] A. Lyaghfouri : A Regularity Result for a Heterogeneous Evolution Dam Problem. Zeitschrift für Analysis und ihre Anwendungen, Vol. 24, No. 1, 149-166 (2005).

[14] A. Lyaghfouri, E. Zaouche: *L^p-continuity of solutions to parabolic free boundary problems*. Electronic Journal of Differential Equations, Vol. 2015, No. 184, pp. 1-9 (2015).

[15] A. Torelli, *Existence and uniqueness of the solution of a non steady free boundary problem*. Boll. U.M.I., 14-B(5), 423-466 (1977).

[16] E. Zaouche: *Existence of a Solution in a Class of Parabolic Free Boundary Problems*. Submitted.